

1/7

DIFFERENCES IN THE MEANS OF THE INITIAL
SChE LEVELS IN THE FOUR STUDY GROUPS




FIG. 1




FIG. 2

CORRELATION OF VAS WITH SChE:
AS THE VAS RISES, THE PERCENT OF CASES WITH
PREOPERATIVE SChE LEVELS ABOVE 1272 TEND TO RISE




FIG. 3A





FIG. 3B


2/7


PREOPERATIVE AND 30 MINUTES
POST-ANESTHETIC INDUCTION
SCHe LEVELS IN CSP PATIENTS
AND SURGICAL CONTROLS



PREOPERATIVE TO 30 MINUTES
POST-INDUCTION SCHe LEVELS IN 55 CSP
PATIENTS, 11 SURGICAL CONTROLS, AND
IN 13 CSP PATIENTS SAMPLED 1 DAY PO

FIG. 3C






3/7

FIG. 3D

MEANS OF THE PREOPERATIVE AND 30 MINUTES POST-INDUCTION SChe
LEVELS OF SEVEN CSP PATIENTS IN WHICH THE LEVEL ROSE > 1 SD
THE VOLATILITY OF CASE 25 IS ILLUSTRATED



MEAN SChe LEVELS OF NINE DISABLED
PATIENTS AND EIGHT CSP PATIENTS






FIG. 4




4/7

FIG. 5

PREOPERATIVE TO 30 MINUTES POST-INDUCTION SCHE LEVELS IN 54 CSP PATIENTS






5/7

FIG. 6

PREOPERATIVE TO 30 MINUTES POST-INDUCTION SChe LEVELS IN 34 CSP PATIENTS WITH PREOPERATIVE LEVELS > 1272







6/7

FIG. 7

PREOPERATIVE TO 30 MINUTES POST-INDUCTION SCHE LEVELS IN 20 CSP PATIENTS WITH PREOPERATIVE LEVELS < 1272*





7/7

FIG. 8

PRE TO INTRAOPERATIVE SCHe LEVELS IN 11 INSTRUMENTED CSP CASES THAT WERE MONITORED

